




Field Scabious

Knautia arvensis Bachelor's buttons, Gypsy rose

Provincial Designation:
Noxious



Alberta Sustainable Resource Development

Overview:

Field scabious is native to Europe and was introduced as an ornamental plant. It is a tall perennial that favors grassy areas and develops a deep tap root. The flowers very closely resemble those of another ornamental perennial, *Scabiosa* (butterfly plant, pincushion flower) but each plant belongs to a different genus.

Habitat:

Prefers nutrient-rich and moderately dry soils, but can also establish in gravelly soils.

Identification:

Stems: Are erect, hairy, sparsely branched, and grow up to 1.5 m tall. There can be one or several stems per plant, with little or no branching in the upper stem.

Leaves: Are hairy and the degree of the lobes is highly variable. Young rosettes leaves tend to be lance-shaped, have pointed tips, and the margins can be entire or coarsely toothed – sometimes a few leaves will be

pinnately lobed. Stem leaves are opposite, pinnate (deeply lobed) and attached directly to the stem. Lower leaves are 10-25 cm long but become smaller higher on the plant.

Flowers: Are a composite of small, violet-blue to purple florets clustered into a head resembling a single flower up to 4 cm wide, and occur singly at the ends of stems. Occasionally flowering stems arise from leaf axils lower on the stem. Below the flower head is a ring of narrow green bracts. Flowers are hermaphroditic (having both male and female organs).

Seeds: Once flowering is complete the seed head is domed and covered with short, bristly hairs. The fruit is nut like, cylindrical and hairy, 5-6mm in size. Seeds fall around the parent plant. A single plant can produce up to 2000 seeds that remain viable for many years.

Prevention:

Field scabious can invade undisturbed plant communities, and once established is very difficult to control, but maintaining healthy cover can help to prevent against invasion.

Any field scabious infestation that has been allowed to go to seed a few times will require many years of diligent control work to eradicate. Since this plant favors grassy areas such as hayfields, it can be widely dispersed in baled forage.

Control:

Grazing: Field scabious is not palatable and seeds can be transported by animal movement. Invasive plants should never be considered as forage.

Cultivation: Discing before flowering is effective in crop land situations.

Mechanical: Mowing is effective to prevent seed production but would likely need to be repeated in the season because of re-sprouting. The deep tap root is difficult to remove in anything but loose soils; therefore hand pulling usually results in the stem breaking off at ground level and then re-sprouting occurs. However, any removal of seed is beneficial. Wear long sleeves and gloves as skin contact with the hairy plant causes considerable itching.

continued next page

Field Scabious (Continued)

Chemical: Metsulfuron-methyl alone or in a product mix with Aminopyralid are registered for use on field scabious. Always check product labels to ensure the herbicide is registered for use on the target plant in Canada by the Pest Management Regulatory Agency. Always read and follow label directions. Consult your local Agricultural Fieldman or Certified Pesticide Dispenser for more information.

Biological: None researched to date.



Alberta Sustainable Resource Development



Alberta Sustainable Resource Development



Alberta Sustainable Resource Development